

Continued fractions with minimal remainders

Jabitskaya E. N.¹

Consider the representation of a rational number in the form, associated with "centered" Euclidean algorithm. We prove a new formula for the limit distribution function for sequences of rationals with bounded sum of partial quotients.

1 Introduction and main results

The classical Euclidean algorithm for $a \in \mathbb{Z}$ and $b \in \mathbb{N}$ uses the division of the form

$$a = bq + r, \quad q \in \mathbb{Z}, \quad b > 0, \quad 0 \leq r < b.$$

It leads to a continued fraction expansion of a real number

$$x = [a_0; a_1, a_2, \dots, a_m, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots + \frac{1}{a_m + \dots}}}, \quad (1)$$

where $a_0 \in \mathbb{Z}$, $a_j \in \mathbb{N}$ for $j \geq 1$. Numbers a_i are called partial quotients of fraction (1).

For $x \in \mathbb{Q}$ the representation (1) is finite. We assume for the uniqueness that the last partial quotient a_l is greater or equal than 2. Let

$$S^{[0]}(a/b) := a_0 + a_1 + \dots + a_l.$$

Define the set

$$\mathcal{F}_n := \{x \in \mathbb{Q}, x \in [0, 1] : S^{[0]}(x) \leq n + 1\}. \quad (2)$$

The limit distribution function

$$F^{[0]}(x) := \lim_{n \rightarrow \infty} \frac{\#\{\xi \in \mathcal{F}_n : \xi \leq x\}}{\#\mathcal{F}_n}, \quad x \in [0, 1]$$

is the famous Minkowski's question mark function $?(x)$. Properties of $?(x)$ were investigated for example in [1], [14] and [2].

¹ Research is supported by RFBR grant No 09-01-00371-a

There are different kinds of Euclidean algorithms. For example, "by-excess" Euclidean algorithm uses the division "by-excess"

$$a = bq + r, \quad -b < r \leq 0,$$

This algorithm leads to regular reduced continued fraction ([3], [5]) expansion of a real number x , that is

$$x = [[a_0; a_1, a_2, \dots, a_m, \dots]] = a_0 - \frac{1}{a_1 - \frac{1}{a_2 - \dots - \frac{1}{a_m - \dots}}}, \quad (3)$$

where $a_0 \in \mathbb{Z}$, $a_j \geq 2$ for $j \geq 1$. Numbers a_i are called partial quotients of fraction (3).

For a rational x representation (3) is finite. For rational x we denote the sum of partial quotients in the representation of x in the form (3) by $S^{[1]}(x)$. We put

$$\Xi_n := \{x \in \mathbb{Q}, x \in [0, 1] : S^{[1]}(x) \leq n + 2\}$$

Consider the limit distribution function

$$F^{[1]}(x) := \lim_{n \rightarrow \infty} \frac{\#\{\xi \in \Xi_n : \xi \leq x\}}{\#\Xi_n}, \quad x \in [0, 1],$$

In 1995 R. F. Tichy and J. Uitz [8] considered a one parameter family $g_\lambda(x)$, $\lambda \in (0, 1)$, $x \in [0, 1]$, of singular functions. Functions $F^{[0]}$, $F^{[1]}$ belong to this family with $\lambda = \frac{1}{2}$ and $\lambda = \frac{3-\sqrt{5}}{2}$ correspondingly. Similar functions $\kappa(x, \alpha)$, $x \in [0, \infty)$, $\lambda \in (0, 1)$, were introduced by A. Denjoy [1] much more earlier in 1938. For $x \in [0, 1]$ functions $\kappa(x, \alpha)$ and $g_\lambda(x)$ are related in the following way:

$$\kappa(x, \alpha) = 1 - (1 - \alpha)g_{1-\alpha}(x).$$

In the same paper A. Denjoy proved that

$$\kappa(x, \alpha) = \alpha^{a_0} - \alpha^{a_0}(1 - \alpha)^{a_1} + \alpha^{a_0+a_2}(1 - \alpha)^{a_1+a_3} - \dots,$$

where $a_0, a_1, \dots, a_m, \dots$ are partial quotients of representation (1) of number x . Similar formula for $g_\lambda(x)$, $x \in (0, 1)$ is given in the paper [5]:

$$g_\lambda(x) = \lambda^{a_1-1} - \lambda^{a_1-1}(1 - \lambda)^{a_2} + \lambda^{a_1-1}(1 - \lambda)^{a_2}\lambda^{a_3} - \dots + \\ + (-1)^{m+1}\lambda^{\sum_{1 \leq i \leq m \atop i \equiv 1 \pmod 2} a_i - 1} (1 - \lambda)^{\sum_{1 \leq i \leq m \atop i \equiv 0 \pmod 2} a_i} + \dots \quad (4)$$

For $\lambda = \frac{1}{2}$ formula (4) gives a well-known result by R. Salem [7], namely

$$g_{1/2}(x) = ?(x) = \frac{1}{2^{a_1-1}} - \frac{1}{2^{a_1+a_2-1}} + \frac{1}{2^{a_1+a_2+a_3-1}} - \dots$$

Let us consider a "centered" version of the Euclidean algorithm. This algorithm uses "centered" division

$$a = bq + r, \quad -\frac{b}{2} < r \leq \frac{b}{2} \quad (5)$$

and leads to the following representation of a real number x :

$$x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l}, \dots \right] = a_0 + \frac{\varepsilon_1}{a_1 + \frac{\varepsilon_2}{a_2 + \dots + \frac{\varepsilon_l}{a_l + \dots}}}. \quad (6)$$

This representation is known as a continued fraction with minimal remainders. Numbers a_i are called partial quotients of fraction (6). Here $a_0 \in \mathbb{Z}$, $\varepsilon_i = \pm 1$ and $a_j \geq 2$, $a_j + \varepsilon_{j+1} > 2$ for $j \geq 1$. For rational x , if $a_s = 2$ is the last partial quotient, then $\varepsilon_s = 1$ for uniqueness of the representation. Such fractions can be found in the book [6] by O. Perron.

Statistical properties of various Euclidean algorithms were investigated by B. Vallee and V. Baladi in papers [11], [12], [13]. The most precise asymptotic formulae for the mean length for the classical Euclidean algorithm and the centered Euclidean algorithm are proved in papers [9], [10] by A. V. Ustinov. A similar formula for "by-excess" Euclidean algorithm was obtained in author's paper [5].

For rational x let us denote by $S(x)$ the sum of partial quotients of representation (6), and put

$$\mathcal{Z}_n := \{x \in \mathbb{Q} \cap [0, 1] : S(x) \leq n + 1\}.$$

In present paper we investigate the limit distribution function

$$F(x) := \lim_{n \rightarrow \infty} \frac{\#\{\xi \in \mathcal{Z}_n : \xi \leq x\}}{\#\mathcal{Z}_n}, \quad x \in [0, 1]. \quad (7)$$

The main result is the following theorem.

Theorem 1. *Let $x \in [0, 1]$, x is represented in the form (6), then*

$$F^{[2]}(x) = a_0 - c\lambda \left(\frac{E_1}{\lambda^{A_1}} + \frac{E_2}{\lambda^{A_2}} + \dots + \frac{E_j}{\lambda^{A_j}} + \dots \right), \quad (8)$$

where

$$E_j = \prod_{1 \leq i \leq j} (-\varepsilon_i), \quad A_j = \sum_{0 \leq i \leq j} a_i, \quad c = 1/(\lambda - 1),$$

and λ is the unique real root of the equation

$$\lambda^3 - \lambda^2 - \lambda - 1 = 0.$$

For rational x the sum in formula (8) is finite.

In this paper we also prove

Theorem 2. *Let for $x \in [0, 1]$ the derivative $F'(x)$ (finite or infinite) exists. Then either $F'(x) = 0$ or $F'(x) = \infty$.*

As function $F(x)$ is monotonic, then by Lebesgue's theorem, the derivative $F'(x)$ exists and is finite almost everywhere (in the sense of Lebesgue measure). That is why $F'(x) = 0$ almost everywhere. In other words, $F(x)$ is a singular function.

In the proof of Theorem 1 we need the following result.

Proposition 1. *For $x \in [0, 1/2]$ function $F(x)$ satisfies the following functional equation*

$$F(1 - x) = 1 - \frac{F(x)}{\lambda}. \quad (9)$$

The proof of Proposition 1 is given in section 3. Theorem 1 uses Proposition 1 and its proof is given in section 4. The proof of Theorem 2 is in section 5.

2 Properties of a continued fraction with minimal remainders

It follows immediately from the definition of continued fraction with minimal remainders (6), that

- $a_i \geq 2$ for $i \geq 1$,
- If $a_i = 2$, then $\varepsilon_{i+1} = 1$ for $i \geq 1$,
- If the last partial quotient $a_l = 2$, then $\varepsilon_l = 1$.

Let $x = [b_0; b_1, \dots, b_s, \dots]$ is represented in the form of ordinary continued fraction (1). We describe the algorithm for converting this fraction into a fraction of the form (6) (see [6]).

Fraction (1) is constructed by the classical Euclidean algorithm

$$r_0 = \frac{b}{a}, \quad r_{i+1} = \frac{1}{r_i} - b_i, \quad b_i = \left[\frac{1}{r_i} \right], \quad 0 \leq r_i < 1.$$

The remainder r_{i+1} is less than $\frac{1}{2}$ if and only if $b_{i+1} > 1$. So while $b_{i+1} > 1$ partial quotients of the classical Euclidean algorithm coincide with partial quotients of "centered" Euclidean algorithm.

For the first i such that $b_{i+1} = 1$, we use the identity

$$b_i + \frac{1}{1 + \frac{1}{b_{i+2} + \alpha}} = b_i + 1 - \frac{1}{b_{i+2} + 1 + \alpha}, \quad \alpha \geq 0.$$

And since $\frac{1}{b_{i+1} + 1} < \frac{1}{2}$, we have

$$[b_0; b_1, \dots, b_i, 1, b_{i+2}] = \left[b_0; \frac{1}{b_1}, \dots, \frac{1}{b_i + 1}, \frac{-1}{b_{i+2} + 1} \right]$$

Then we apply the same procedure to the "tail"

$$b_{i+2} + 1 + \frac{1}{b_{i+3} + \dots + \frac{1}{b_s + \dots}}$$

of the fraction (1).

We define the convergents of the continued fraction with minimal remainders of the number $x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l}, \dots \right]$ as

$$\frac{P_n}{Q_n} = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_n}{a_n} \right], \quad (P_n, Q_n) = 1, \quad n \geq 0.$$

To get a recurrence formulas for P_n/Q_n , $n \geq 0$ we put formally

$$\frac{P_{-1}}{Q_{-1}} = \frac{1}{0}, \quad \frac{P_0}{Q_0} = \frac{a_0}{1}.$$

Then for $\varepsilon_{n+1} = 1$ we have

$$\frac{P_{n+1}}{Q_{n+1}} = \frac{a_n P_n + P_{n-1}}{a_n Q_n + Q_{n-1}},$$

otherwise

$$\frac{P_{n+1}}{Q_{n+1}} = \frac{a_n P_n - P_{n-1}}{a_n Q_n - Q_{n-1}}.$$

3 Definition and properties of sets $\mathcal{Z}_n$

We define a sequence of sets $\mathcal{X}_k$ by

$$\mathcal{X}_k = \{x \in \mathbb{Q} \cap [0, 1] : S(x) = k + 1\}, \quad n \geq 1.$$

It is clear that

$$\mathcal{Z}_n = \bigcup_{1 \leq k \leq n} \mathcal{X}_k,$$

where $\mathcal{Z}_n$ is defined by (7). Suppose that the elements of $\mathcal{Z}_k$ are arranged in the increasing order. The number of elements of $\mathcal{Z}_n$, $\mathcal{X}_n$ we denote by Z_n , X_n correspondingly.

$$\text{Particularly, } \mathcal{X}_1 = \left\{ \frac{1}{2} \right\}, \mathcal{X}_2 = \left\{ \frac{1}{3} \right\}, \mathcal{X}_3 = \left\{ \frac{1}{4}, \frac{2}{5}, \frac{2}{3} \right\}, \mathcal{X}_4 = \left\{ \frac{1}{5}, \frac{2}{7}, \frac{3}{7}, \frac{3}{5}, \frac{4}{4} \right\}.$$

So $X_1 = X_2 = 1$, $X_3 = 3$, $X_4 = 5$.

Lemma 1. *For $n \geq 1$ we have*

$$X_{n+3} = X_{n+2} + X_{n+1} + X_n.$$

Proof. We construct one-to-one correspondence Φ between elements of sets $\mathcal{X}_{n+2} \cup \mathcal{X}_{n+1} \cup \mathcal{X}_n$ and $\mathcal{X}_{n+3}$.

Let $x \in \mathcal{X}_{n+2} \cup \mathcal{X}_{n+1} \cup \mathcal{X}_n$, $x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l} \right]$, we define $\Phi(x) : \mathcal{X}_{n+2} \cup \mathcal{X}_{n+1} \cup \mathcal{X}_n \rightarrow \mathcal{X}_{n+3}$ in the following way:

- If $x \in \mathcal{X}_{n+2}$, then

$$\Phi(x) = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}}, \frac{\varepsilon_l}{a_l + 1} \right] \in \mathcal{X}_{n+3}.$$

- If $x \in \mathcal{X}_{n+1}$, then

$$\Phi(x) = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l}, \frac{1}{2} \right] \in \mathcal{X}_{n+3}.$$

- If $x \in \mathcal{X}_n$ and $a_l > 2$, then

$$\Phi(x) = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l}, \frac{-1}{3} \right] \in \mathcal{X}_{n+3}.$$

- If $x \in \mathcal{X}_n$ and $a_l = 2$, then

$$\Phi(x) = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-2}}{a_{l-2}}, \frac{\varepsilon_{l-1}}{a_{l-1} + 1}, \frac{-1}{2}, \frac{1}{2} \right] \in \mathcal{X}_{n+3}.$$

The correspondence $\Phi(x)$ is injective by the construction. Let us show that it is surjective. For any $y \in \mathcal{X}_{n+3}$, $y = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l} \right]$ we find the preimage x of y .

- If $a_l > 3$ or $a_l = 3$ and $\varepsilon_l = 1$ then

$$x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}}, \frac{\varepsilon_l}{a_l - 1} \right] \in \mathcal{X}_{n+2}.$$

- If $a_l = 2$ and either $a_{l-1} > 2$ or $a_{l-1} = 2$ and $\varepsilon_{l-1} = 1$, then

$$x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}} \right] \in \mathcal{X}_{n+1}.$$

- If $a_l = 3$, $\varepsilon_l = -1$, then $a_{l-1} > 2$, therefore

$$x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}} \right] \in \mathcal{X}_n.$$

- If $a_l = a_{l-1} = 2$, $\varepsilon_{l-1} = -1$, then $a_{l-2} > 2$, therefore

$$x = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-3}}{a_{l-3}}, \frac{\varepsilon_{l-2}}{a_{l-2}-1}, \frac{1}{2} \right] \in \mathcal{X}_n.$$

Lemma is proved. ■

Corollary 1. *For $n \geq 1$ we have*

$$Z_{n+3} = Z_{n+2} + Z_{n+1} + Z_n + 2. \quad (10)$$

Proof. By the definition of Z_n and Lemma 1, we get

$$\begin{aligned} Z_{n+2} + Z_{n+1} + Z_n &= \\ &= (X_1 + \dots + X_{n+2}) + (X_1 + \dots + X_{n+1}) + (X_1 + \dots + X_n) = \\ &= X_1 + X_2 + X_3 + X_4 + \dots + X_{n+3} + (X_1 - X_3) = Z_{n+3} - 2. \end{aligned}$$

■

We remind the definition of the Stern-Brocot sequences $\mathcal{F}_n$, $n = 0, 1, 2, \dots$. Consider two-point set $\mathcal{F}_0 = \left\{ \frac{0}{1}, \frac{1}{1} \right\}$. Let $n \geq 0$ and

$$\mathcal{F}_n = \{0 = x_{0,n} < x_{1,n} < \dots < x_{N(n),n} = 1\},$$

where $x_{j,n} = p_{j,n}/q_{j,n}$, $(p_{j,n}, q_{j,n}) = 1$, $j = 0, \dots, N(n)$ and $N(n) = 2^n + 1$. Then

$$\mathcal{F}_{n+1} = \mathcal{F}_n \cup Q_{n+1}$$

with

$$Q_{n+1} = \{x_{j-1,n} \oplus x_{j,n}, \quad j = 1, \dots, N(n)\}.$$

Here

$$\frac{a}{b} \oplus \frac{c}{d} = \frac{a+b}{c+d}$$

is the mediant of fractions $\frac{a}{b}$ and $\frac{c}{d}$.

So the first sequences are $Q_1 = \left\{ \frac{1}{2} \right\}$, $Q_2 = \left\{ \frac{1}{3}, \frac{2}{3} \right\}$, $Q_3 = \left\{ \frac{1}{4}, \frac{2}{5}, \frac{3}{5}, \frac{3}{4} \right\}$.

It is clear that for any rational number q there exists such number n that $q \in Q_n$. Note that sum $S^{[0]}(x)$ of partial quotients of ordinary continued fraction of a number $x \in Q_n$ equals to $n+1$. Formula (2) gives an equivalent definition of $\mathcal{F}_n$.

It is convenient to represent sequences $\mathcal{F}_n$ by means of the binary tree $\mathcal{D}^{[0]}$ (Figure 1). This tree is called Stern-Brocot's tree. Nodes of the




Figure 1:

tree are labeled by rationals from $(0,1)$ and partitioned into levels by the following rule: n -th level consists of nodes labeled by numbers x , such that $S^{[0]}(x) = n+1$ (i.e. n -th level consists of nodes, labeled by numbers from Q_n).

It is possible to distribute nodes of the tree into levels by another way. For example, we can use such a rule: n -th level consists of nodes labeled by numbers x , such that sum $S^{[1]}(x)$ of partial quotients of regular reduced continued fraction of number x equals $n+1$. Then we get tree $\mathcal{D}^{[1]}$ (Figure 2) from paper [4].

Now let us distribute nodes of the tree into levels by the following rule: n -th level consists of nodes labeled by numbers x , such that $S(x) = n+1$ (i.e. $x \in \mathcal{X}_n$). We denote this tree by $\mathcal{D}$ (Figure 3).

Any node ξ of the tree $\mathcal{D}$ is a root of subtree, which we denote by $\mathcal{D}^{(\xi)}$. Nodes of $\mathcal{D}^{(\xi)}$ are also partitioned into levels: ξ itself belongs to level 1, and a node of the tree $\mathcal{D}^{(\xi)}$, labeled by number x belongs to the level $S(x) - S(\xi) + 1$




Figure 2:

in the tree $\mathcal{D}^{(\xi)}$. The number of nodes of $\mathcal{D}^{(\xi)}$ from level 1 to level n we denote by $D_n^{(\xi)}$.

Let us consider more detailed structure of the tree $\mathcal{D}$. From every node ξ of $\mathcal{D}$ we issue two arrows: the left one and the right one. The left one goes to the node labeled by x^l and the right one goes to node labeled by x^r . Note that if $\xi = x \oplus y$, where x, y are consecutive elements of $\mathcal{F}_n$, then $\xi^l = x \oplus \xi$, $\xi^r = \xi \oplus y$.

Remark 1. Let $\xi \in \mathcal{X}_n$ and $\xi = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l} \right] = x \oplus y$, where x, y — neighboring elements of $\mathcal{F}_n$, $S(x) < S(y)$. If $a_l > 2$, then

$$x \oplus \xi = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}}, \frac{\varepsilon_l}{a_l + 1} \right] \in X_{n+1},$$

$$y \oplus \xi = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}}, \frac{\varepsilon_l}{a_l - 1}, \frac{1}{2} \right] \in X_{n+1},$$

If $a_l = 2$, then $\varepsilon_l = 1$ and

$$x \oplus \xi = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}}, \frac{1}{3} \right] \in X_{n+1},$$




Figure 3:

$$y \oplus \xi = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_{l-1}}{a_{l-1}}, \frac{\varepsilon_{l-1}}{a_{l-1} + 1}, \frac{-1}{3} \right] \in X_{n+2}.$$

From Remark 1 we deduce the following statement.

Lemma 2. *Let $\xi = \left[a_0; \frac{\varepsilon_1}{a_1}, \dots, \frac{\varepsilon_l}{a_l} \right]$, then*

$$D_n^{(\xi)} = \begin{cases} D_n^{(1/2)}, & \text{if } a_l = 2 \\ D_n^{(1/3)}, & \text{if } a_l > 2. \end{cases} \quad (11)$$

Note that $D_n^{(1/2)} = Z_n$. For brevity we put $Y_n = D_n^{(1/3)}$. So $Y_1 = 1$, $Y_2 = 2$, $Y_3 = 3$.

For Z_n we have recurrence formula (10). It is easy to prove a similar formula for Y_n :

$$Y_{n+3} = Y_{n+2} + Y_{n+1} + Y_n + 2. \quad (12)$$

Lemma 3. *Let λ be the unique real root of the equation*

$$\lambda^3 - \lambda^2 - \lambda - 1 = 0, \quad (13)$$

and $c = 1/(\lambda - 1)$. Then

$$\lim_{n \rightarrow \infty} \frac{Y_n}{Y_{n+1}} = \lim_{n \rightarrow \infty} \frac{Z_n}{Z_{n+1}} = \lambda, \quad \lim_{n \rightarrow \infty} \frac{Y_n}{Z_n} = c.$$

Proof. Equation (12) can be reduced to a homogeneous by the substitution $Y'_n = Y_n + 1$:

$$Y'_{n+3} = Y'_{n+2} + Y'_{n+1} + Y'_n.$$

The characteristic equation is

$$\lambda^3 - \lambda^2 - \lambda = 1.$$

This equation has the unique real root $\lambda \approx 1,839292$ and two complex roots λ_1, λ_2 , such that $|\lambda_1| = |\lambda_2| < 1$. So

$$\begin{aligned} Y_n + 1 = Y'_n &= C_1 \lambda^n + C_2 \lambda_2^n + C_3 \lambda_3^n, \\ Z_n + 1 = Z'_n &= D_1 \lambda^n + D_2 \lambda_2^n + D_3 \lambda_3^n, \end{aligned}$$

with certain constants $C_1, C_2, C_3, D_1, D_2, D_3$. Put $c = C_1/D_1$. From construction of the tree $\mathcal{D}$ it is clear that

$$Z_n = Y_{n-1} + Y_{n-2} + 1.$$

Dividing both parts of this equality by Z_n and taking the limit we get

$$\begin{aligned} 1 &= \lim_{n \rightarrow \infty} \frac{C_1 (\lambda_1^{n-1} + \lambda_1^{n-2}) + C_2 (\lambda_2^{n-1} + \lambda_2^{n-2}) + C_3 (\lambda_3^{n-1} + \lambda_3^{n-2}) + 1}{D_1 \lambda_1^n + D_2 \lambda_2^n + D_3 \lambda_3^n + 1} = \\ &= \frac{C_1 + C_1 \lambda}{D_1 \lambda^2}. \end{aligned}$$

Since λ is the root of equation (13), we get the following relation between c and λ :

$$c = \frac{\lambda^2}{1 + \lambda} = \frac{1}{\lambda - 1} \approx 1,1915. \quad (14)$$

Lemma is proved. ■

4 Properties of the limit distribution function $F(x)$ of sequence $\mathcal{Z}_n$

In this section we prove some auxiliary results about function $F(x)$.

Lemma 4. *Let x, y be consecutive elements of the sequence $\mathcal{Z}_n$. Then*

$$F(y) - F(x) = \lim_{n \rightarrow \infty} \frac{D_{n+2-S(x \oplus y)}^{(x \oplus y)}}{D_n^{(1/2)}}.$$

Proof. Note that nodes of tree $\mathcal{D}^{(x \oplus y)}$ are labeled exactly by the numbers from the set $\{\xi \in \mathbb{Q} : x < \xi < y\}$. So

$$F(y) - F(x) = \lim_{n \rightarrow \infty} \frac{\#\{\xi \in \mathcal{Z}_n : x < \xi \leq y\}}{Z_n} = \lim_{n \rightarrow \infty} \frac{D_{n+2-S(x \oplus y)}^{(x \oplus y)}}{D_n^{(1/2)}}.$$

■

Lemma 5. *Let x, y be consecutive elements of $\mathcal{Z}_n$, $S(x) < S(y)$, and let a_l be the last partial quotient in continued fraction with minimal remainders representation of the number $x \oplus y$.*

If $a_l = 2$, then

$$F(x \oplus y) - F(x) = \frac{c}{\lambda} (F(y) - F(x)), \quad (15)$$

$$F(x \oplus y) - F(y) = \frac{c}{\lambda^2} (F(x) - F(y)). \quad (16)$$

If $a_l > 2$, then

$$F(x \oplus y) - F(x) = \frac{1}{\lambda} (F(y) - F(x)), \quad (17)$$

$$F(x \oplus y) - F(y) = \frac{1}{c\lambda} (F(x) - F(y)). \quad (18)$$

Proof. We suppose that $x < y$ (in case $x > y$ the proof is similar). According to Lemma 4 one has

$$F(x \oplus y) - F(x) = \lim_{n \rightarrow \infty} \frac{D_{n+2-S((x \oplus y)^l)}^{((x \oplus y)^l)}}{D_n}$$

$$F(y) - F(x \oplus y) = \lim_{n \rightarrow \infty} \frac{D_{n+2-S((x \oplus y)^r)}^{((x \oplus y)^r)}}{D_n}$$

By Remark 1, if $a_l = 2$, then the last partial quotients of continued fractions with minimal remainders of numbers $(x \oplus y)^l$, $(x \oplus y)^r$ are greater then 2 and

$$S((x \oplus y)^l) = S(x \oplus y) + 1, \quad S((x \oplus y)^r) = S(x \oplus y) + 2.$$

So

$$\frac{F(x \oplus y) - F(x)}{F(y) - F(x \oplus y)} = \lim_{n \rightarrow \infty} \frac{D_{n+2-S((x \oplus y)^l)}^{((x \oplus y)^l)}}{D_{n+2-S((x \oplus y)^r)}^{((x \oplus y)^r)}} = \lim_{n \rightarrow \infty} \frac{Y_{n+1-S(x \oplus y)}}{Y_{n-S(x \oplus y)}} = \lambda. \quad (19)$$

i.e. $F(x \oplus y)$ divides the segment $[F(x), F(y)]$ in the relation $\lambda : 1$. Taking into account that

$$\frac{c}{\lambda^2} + \frac{c}{\lambda} = 1$$

we get formulas (15), (16).

If $a_l > 2$, then by Remark 1 the last partial quotient of the continued fraction with minimal remainders for the number $(x \oplus y)^r$ is 2. For the number $(x \oplus y)^l$ the last partial quotient is greater then 2 and

$$S((x \oplus y)^l) = S((x \oplus y)^r) = S(x \oplus y) + 1.$$

That is why

$$\frac{F(x \oplus y) - F(x)}{F(y) - F(x \oplus y)} = \lim_{n \rightarrow \infty} \frac{D_{n+2-S(x \oplus y)^l}^{((x \oplus y)^l)}}{D_{n+2-S(x \oplus y)^r}^{((x \oplus y)^r)}} = \lim_{n \rightarrow \infty} \frac{Y_{n+1-S(x \oplus y)}}{Z_{n+1-S(x \oplus y)}} = c. \quad (20)$$

i.e. the number $F(x \oplus y)$ divides the segment $[F(x), F(y)]$ in the relation $c : 1$. Taking into account that

$$\frac{1}{\lambda} + \frac{1}{c\lambda} = 1$$

we get formulas (17), (18). Lemma is proved. ■

Now we are able to prove Proposition 1. Suppose $x \in [0, 1/2]$. By the definition

$$F(x) = \lim_{n \rightarrow \infty} \frac{\#\{\xi \in \mathcal{Z}_n : \xi \leq x\}}{Z_n}.$$

So

$$\lim_{n \rightarrow \infty} \frac{\# \{ \xi \in \mathcal{Z}_{n-1} : \xi \leq x \}}{Z_{n-1}} \frac{Z_{n-1}}{Z_n} = \frac{F^{[2]}(x)}{\lambda}.$$

Taking into account that $S(1 - \xi) = 1 + S(\xi)$ for $\xi \in \mathbb{Q} \cap [0, 1/2]$, for $x \in [0, 1/2]$ we have

$$\begin{aligned} \{ \xi \in \mathcal{Z}_{n-1} : \xi < x \} &= \{ 1 - \xi = \eta \in \mathcal{Z}_n : 1 - \eta < x \} = \\ &= \{ \eta \in \mathcal{Z}_n : \eta > 1 - x \}. \end{aligned}$$

So we get

$$\begin{aligned} \frac{F^{[2]}(x)}{\lambda} + F^{[2]}(1 - x) &= \\ &= \lim_{n \rightarrow \infty} \frac{\# \{ \xi \in \mathcal{Z}_{n-1} : \xi \leq x \}}{Z_n} + \lim_{n \rightarrow \infty} \frac{\# \{ \xi \in \mathcal{Z}_n : \xi \leq 1 - x \}}{Z_n} = \\ &= \lim_{n \rightarrow \infty} \frac{\# \{ \eta \in \mathcal{Z}_n : \eta > 1 - x \} \cup \{ \xi \in \mathcal{Z}_n : \xi \leq 1 - x \}}{Z_n} = 1. \end{aligned}$$

This equality proofs formula (9). ■

5 Proof of Theorem 1

Let us prove the theorem for rational $x \in [0, 1/2]$ by induction on $S(x)$. The equality

$$F(1/a_1) = \frac{c}{\lambda^{a_1-1}},$$

follows from formula (15), since $1/a_1 = \underbrace{0 \oplus \dots \oplus 0}_{(a_1-1) \text{ times}} \oplus 1$.

Suppose that the formula (8) is proved for

$$x = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m} \right].$$

Then it is enough to prove it for

$$y = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{a_m + 1} \right]$$

and for

$$z = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m - 1}, \frac{1}{2}\right], \quad \text{if } a_m > 2,$$

$$w = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1} + 1}, \frac{-1}{3}\right], \quad \text{if } a_m = 2.$$

We see that

$$y = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}\right] \oplus \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{a_m}\right]$$

and the last partial quotient $a_m + 1$ of continued fraction with minimal remainders expression of number y is greater then 2. From (17) and the inductive assumption we get

$$\begin{aligned} F(y) &= F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}\right]\right) + \\ &+ \frac{1}{\lambda} \left(F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m}\right]\right) - F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}\right]\right) \right) = \\ &= F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}\right]\right) - \frac{1}{\lambda} c \lambda \frac{E_m}{\lambda^{A_m}} = \\ &= F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}\right]\right) - c \lambda \frac{E_m}{\lambda^{A_{m-1} + (a_m + 1)}}. \end{aligned}$$

If $a_m > 2$, we must prove the formula (8) for z . We see that

$$z = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m}\right] \oplus \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{a_m - 1}\right]$$

and the last partial quotient of number z is 2. So by (15) we have

$$\begin{aligned} F(z) &= F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{a_m - 1}\right]\right) - \\ &- \frac{c}{\lambda} \left(F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{a_m - 1}\right]\right) - F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m}\right]\right) \right) = \\ &= F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{1}{a_m}\right]\right) + \frac{c}{\lambda} c \lambda \left(\frac{E_m}{\lambda^{A_{m-1} + (a_m - 1)}} - \frac{E_m}{\lambda^{A_m}} \right) = \\ &= F\left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{1}{a_m}\right]\right) - c \lambda \frac{E_m(-1)}{\lambda^{A_{m-1} + (a_m - 1) + 2}}. \end{aligned}$$

If $a_m = 2$, we must prove formula (8) for w . We see that

$$w = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{a_m + 1}\right] \oplus \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m}, \frac{1}{2}\right]$$

and the last partial quotient of number w is greater than 2. So by (17) we have

$$\begin{aligned} F^{[2]}(w) &= F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1} + 1}\right] \right) + \\ &\quad + \frac{1}{\lambda} \left(F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{1}{2}\right] \right) - \right. \\ &\quad \left. - F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1} + 1}\right] \right) \right), \quad (21) \end{aligned}$$

As

$$\begin{aligned} \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{1}{2}\right] &= \\ &= \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1} + 1}\right] \oplus \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1}}\right], \end{aligned}$$

by (16) we have

$$\begin{aligned} &F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{1}{2}\right] \right) - \\ &\quad - F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1} + 1}\right] \right) = \\ &\quad = \frac{c}{\lambda^2} \left(F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}\right] \right) - \right. \\ &\quad \left. - F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1} + 1}\right] \right) \right) = \\ &\quad = -\frac{c}{\lambda^2} c\lambda E_{m-1} \left(\frac{1}{\sum_{\lambda^0 \leq i \leq m-2} a_i + a_{m-1}} - \frac{1}{\sum_{\lambda^0 \leq i \leq m-2} a_i + (a_{m-1} + 1)} \right) = \\ &\quad = -c\lambda E_{m-1} \frac{1}{\sum_{\lambda^0 \leq i \leq m-1} a_i + 3}. \end{aligned}$$

Substituting this result in (21), we finally get

$$F^{[2]}(w) = F^{[2]} \left(\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-2}}{a_{m-2}}, \frac{\varepsilon_{m-1}}{a_{m-1} + 1} \right] \right) - c\lambda E_{m-1} \frac{1}{\lambda^{A_{m-1}+4}}.$$

So we have proven Theorem 1 for rational $x \in [0, 1/2]$. For rational $x \in (1/2, 1]$ it follows from formula (9). For irrational $x \in [0, 1]$ we should take into account the continuity of $F(x)$. ■

6 Singularity of the function $F(x)$

In this section we prove Theorem 2. First of all let us consider the case $x \in \mathbb{Q}$.

Lemma 6. *For rational $x \in [0, 1]$ we have $F'(x) = 0$.*

Let us prove that the right derivative $F'_+(x)$ exists and $F'_+(x) = 0$ (for $F'_-(x)$ the prove is similar).

Let $x = a/b$, $a, b \in \mathbb{N}$, then there exists such n , that $a/b \in \mathcal{X}_n$. Denote by a'/b' the right neighbouring to a/b element in $\mathcal{Z}_n$. Sequence of mediant $y_k = \left\{ \frac{ka+a'}{kb+b'} \right\}$, converges to a/b from the right as $k \rightarrow \infty$. So for $\xi > x$ sufficiently close to x there exists such m that $x < y_{m+1} \leq \xi \leq y_m$ and so

$$\frac{|F(\xi) - F(a/b)|}{\xi - a/b} \leq \frac{F(y_m) - F(a/b)}{y_{m+1} - a/b}.$$

Remind that $a/b \in \mathcal{Z}_n$, but $a/b \notin \mathcal{Z}_{n-1}$. So $S(a/b) > S(a'/b')$. By Lemma 5 we see that

$$\begin{aligned} F(a/b \oplus a'/b') - F(a/b) &\leq \max \left(\frac{c}{\lambda^2}, \frac{1}{c\lambda} \right) (F(a'/b') - F(a/b)) = \\ &= \frac{1}{c\lambda} (F(a'/b') - F(a/b)). \end{aligned}$$

Similarly $S \left(\frac{ka+a'}{kb+b'} \right) > S(a/b)$, $k = 1, \dots, m-1$, and

$$\begin{aligned} F \left(\frac{(k+1)a+a'}{(k+1)b+b'} \right) - F(a/b) &\leq \max \left(\frac{c}{\lambda}, \frac{1}{\lambda} \right) \left(F \left(\frac{ka+a'}{kb+b'} \right) - F(a/b) \right) \leq \\ &\leq \frac{c^k}{\lambda^k} (F(a/b \oplus a'/b') - F(a/b)). \end{aligned}$$

So

$$\begin{aligned}
0 \leq F'_+(x) &= \lim_{\xi \rightarrow x_+} \frac{F(\xi) - F(x)}{\xi - x} \leq \lim_{m \rightarrow \infty} \frac{F(y_m) - F(x)}{y_{m+1} - x} = \\
&= \lim_{m \rightarrow \infty} \frac{\frac{c^{m-1}}{\lambda^{m+1}} (F(a'/b') - F(a/b))}{\frac{1}{((m+1)b + b')b}} = 0. \quad (22)
\end{aligned}$$

■

Let $x \notin \mathbb{Q}$. Suppose that $F'(x) = a$, where a is finite and $a \neq 0$. We should prove that it is not possible. We shall use the Stern-Brocot sequences $\mathcal{F}_n$.

Given n we can find two consecutive elements $p_n/q_n < p'_n/q'_n$ from the set $\mathcal{F}_n$ such that $p_n/q_n < x < p'_n/q'_n$. In such way we obtain an infinite sequence of pairs of elements $\{p_n/q_n, p'_n/q'_n\}$, converging to x from the left and from the right correspondingly. So

$$\lim_{n \rightarrow \infty} \frac{F(p'_n/q'_n) - F(p_n/q_n)}{p'_n/q'_n - p_n/q_n} = a,$$

Put

$$G_n(x) = \frac{F(p'_{n+1}/q'_{n+1}) - F(p_{n+1}/q_{n+1})}{F(p'_n/q'_n) - F(p_n/q_n)} \frac{q_{n+1}q'_{n+1}}{q_nq'_n}.$$

Then

$$G_n(x) = \frac{F(p'_{n+1}/q'_{n+1}) - F(p_{n+1}/q_{n+1})}{p'_{n+1}/q'_{n+1} - p_{n+1}/q_{n+1}} \frac{p'_n/q'_n - p_n/q_n}{F(p'_n/q'_n) - F(p_n/q_n)} \xrightarrow{n \rightarrow \infty} 1. \quad (23)$$

It is clear that if $x \in (p_n/q_n, p_n/q_n \oplus p'_n/q'_n)$ then the pair $\{p_{n+1}/q_{n+1}, p'_{n+1}/q'_{n+1}\}$ coincides with $\{p_n/q_n, p_n/q_n \oplus p'_n/q'_n\}$. Also if $x \in (p_n/q_n \oplus p'_n/q'_n, p'_n/q'_n)$ then the pair $\{p_{n+1}/q_{n+1}, p'_{n+1}/q'_{n+1}\}$ coincides with $\{p_n/q_n \oplus p'_n/q'_n, p'_n/q'_n\}$.

Note that $p_n/q_n, p'_n/q'_n$ are always among the intermediate and convergent fractions to x in the sense of ordinary continued fraction.

Let us show that for an irrational x one can find an infinite subsequence $\{p_{n_k}/q_{n_k}, p'_{n_k}/q'_{n_k}\}$ of the sequence $\{p_n/q_n, p'_n/q'_n\}$ with the following property: the last partial quotient in the continued fraction with minimal remainders expression of $p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}$ is equal to 2.

Let

$$x = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_m}{a_m}, \dots \right].$$

Then either

$$p_n/q_n \oplus p'_n/q'_n = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{1}{2}\right],$$

or

$$p_n/q_n \oplus p'_n/q'_n = \left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{b_m}\right],$$

for some natural m , $b_m \leq a_m + 1$. In the first case the pair $\{p_n/q_n, p'_n/q'_n\}$ satisfies the necessary property.

In the other case we consider the pair $\{p_{n+1}/q_{n+1}, p'_{n+1}/q'_{n+1}\}$ and for $p_{n+1}/q_{n+1} \oplus p'_{n+1}/q'_{n+1} \in \{p_{n+2}/q_{n+2}, p'_{n+2}/q'_{n+2}\}$ by remark 1 we have two possibilities: either

$$\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{b_m - 1}, \frac{1}{2}\right]$$

or

$$\left[0; \frac{1}{a_1}, \frac{\varepsilon_2}{a_2}, \dots, \frac{\varepsilon_{m-1}}{a_{m-1}}, \frac{\varepsilon_m}{b_m + 1}\right]$$

where $b_m + 1 \leq a_m + 1$. But the last inequality can occur only finitely many times. So for any m we can find a pair $\{p_{n_k}/q_{n_k}, p'_{n_k}/q'_{n_k}\}$ satisfying the necessary property.

In the sequel we consider such a subsequence $\{p_{n_k}/q_{n_k}, p'_{n_k}/q'_{n_k}\}$. From (23), we see that

$$\lim_{k \rightarrow \infty} G_{n_k}(x) = 1. \quad (24)$$

For a fixed k we consider following cases:

Case 1. $x \in (p_{n_k}/q_{n_k}, p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})$

and

Case 2. $x \in (p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}, p'_{n_k}/q'_{n_k})$.

Consider **Case 1**. As the last partial quotient of $p_n/q_n \oplus p'_n/q'_n$ is equal to 2, Lemma 5 leads to

$$\begin{aligned} \frac{F(p'_{n_k+1}/q'_{n_k+1}) - F(p_{n_k+1}/q_{n_k+1})}{F(p'_{n_k}/q'_{n_k}) - F(p_{n_k}/q_{n_k})} &= \frac{F(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}) - F(p_{n_k}/q_{n_k})}{F(p'_{n_k}/q'_{n_k}) - F(p_{n_k}/q_{n_k})} = \\ &= \begin{cases} \frac{c}{\lambda}, & \text{if } S(p_{n_k}/q_{n_k}) < S(p'_{n_k}/q'_{n_k}). \\ \frac{c}{\lambda^2}, & \text{if } S(p_{n_k}/q_{n_k}) > S(p'_{n_k}/q'_{n_k}). \end{cases} \end{aligned} \quad (25)$$

Then there are two possibilities:

- a) $x \in (p_{n_k}/q_{n_k}, (p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})^l)$,
b) $x \in ((p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})^l, p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})$,
where

$$(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})^l = p_{n_k}/q_{n_k} \oplus (p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}),$$

$$(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})^r = (p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}) \oplus p'_{n_k}/q'_{n_k}.$$

As the last partial quotient of $p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}$ is equal to 2, then by Remark 1, the last partial quotients of $(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})^r$ and $(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})^l$ is not equal to 2. As $S(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}) > S(p_{n_k}/q_{n_k})$, we deduce from Lemma 5 that

$$\frac{F(p'_{n_k+2}/q'_{n_k+2}) - F(p_{n_k+2}/q_{n_k+2})}{F(p'_{n_k+1}/q'_{n_k+1}) - F(p_{n_k+1}/q_{n_k+1})} = \begin{cases} \frac{1}{\lambda}, & \text{in case a),} \\ \frac{1}{c\lambda}, & \text{in case b).} \end{cases} \quad (26)$$

Consider **Case 2**. Analogously to the case 1 we get

$$\begin{aligned} \frac{F(p'_{n_k+1}/q'_{n_k+1}) - F(p_{n_k+1}/q_{n_k+1})}{F(p'_{n_k}/q'_{n_k}) - F(p_{n_k}/q_{n_k})} &= \frac{F(p'_{n_k}/q'_{n_k}) - F(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k})}{F(p'_{n_k}/q'_{n_k}) - F(p_{n_k}/q_{n_k})} = \\ &= \begin{cases} \frac{c}{\lambda^2}, & \text{if } S(p_{n_k}/q_{n_k}) < S(p'_{n_k}/q'_{n_k}). \\ \frac{c}{\lambda}, & \text{if } S(p_{n_k}/q_{n_k}) > S(p'_{n_k}/q'_{n_k}). \end{cases} \end{aligned} \quad (27)$$

We shall consider following subcases:

- a) $x \in (p_n/q_n \oplus p'_n/q'_n, (p_n/q_n \oplus p'_n/q'_n)^r)$,
b) $x \in ((p_n/q_n \oplus p'_n/q'_n)^r, p'_n/q'_n)$.

As $S(p_{n_k}/q_{n_k} \oplus p'_{n_k}/q'_{n_k}) > S(p'_{n_k}/q'_{n_k})$, from Lemma 5 we see that

$$\frac{F(p'_{n_k+2}/q'_{n_k+2}) - F(p_{n_k+2}/q_{n_k+2})}{F(p'_{n_k+1}/q'_{n_k+1}) - F(p_{n_k+1}/q_{n_k+1})} = \begin{cases} \frac{1}{c\lambda}, & \text{in case a),} \\ \frac{1}{\lambda}, & \text{in case b).} \end{cases} \quad (28)$$

As the sequence $\{p_{n_k}/q_{n_k}, p'_{n_k}/q'_{n_k}\}$ is infinite then at least one case (from the cases 1a),1b),2a),2b)) will occur infinitely often. So there exists a subsequence $\{p_{n_{k_m}}/q_{n_{k_m}}, p'_{n_{k_m}}/q'_{n_{k_m}}\}$ such that

$$\frac{F(p'_{n_{k_m}+1}/q'_{n_{k_m}+1}) - F(p_{n_{k_m}+1}/q_{n_{k_m}+1})}{F(p'_{n_{k_m}}/q'_{n_{k_m}}) - F(p_{n_{k_m}}/q_{n_{k_m}})} = \alpha,$$

$$\frac{F(p'_{n_{k_m}+2}/q'_{n_{k_m}+2}) - F(p_{n_{k_m}+2}/q_{n_{k_m}+2})}{F(p'_{n_{k_m}+1}/q'_{n_{k_m}+1}) - F(p_{n_{k_m}+1}/q_{n_{k_m}+1})} = \beta,$$

where α — is one of the numbers $\frac{c}{\lambda}$, $\frac{c}{\lambda^2}$, and β — is one of the numbers $\frac{1}{\lambda}$, $\frac{1}{c\lambda}$. Now

$$G_{n_{k_m}}(x) = \alpha \frac{q_{n_{k_m}+1} q'_{n_{k_m}+1}}{q_{n_{k_m}} q'_{n_{k_m}}}, \quad G_{n_{k_m}+1}(x) = \beta \frac{q_{n_{k_m}+2} q'_{n_{k_m}+2}}{q_{n_{k_m}+1} q'_{n_{k_m}+1}}.$$

From (24) we see that

$$\lim_{m \rightarrow \infty} \frac{q_{n_{k_m}} q'_{n_{k_m}}}{q_{n_{k_m}+1} q'_{n_{k_m}+1}} = \alpha, \quad \lim_{m \rightarrow \infty} \frac{q_{n_{k_m}+1} q'_{n_{k_m}+1}}{q_{n_{k_m}+2} q'_{n_{k_m}+2}} = \beta. \quad (29)$$

Now we must show that (29) is not possible. To do this we distinguish the cases again.

1,a) In this case

$$\begin{aligned} \{p_{n_{k_m}+1}/q_{n_{k_m}+1}, p'_{n_{k_m}+1}/q'_{n_{k_m}+1}\} &= \{p_{n_{k_m}}/q_{n_{k_m}}, (p_{n_{k_m}} + p'_{n_{k_m}})/(q_{n_{k_m}} + q'_{n_{k_m}})\}, \\ \{p_{n_{k_m}+2}/q_{n_{k_m}+2}, p'_{n_{k_m}+2}/q'_{n_{k_m}+2}\} &= \{p_{n_{k_m}}/q_{n_{k_m}}, (2p_{n_{k_m}} + p'_{n_{k_m}})/(2q_{n_{k_m}} + q'_{n_{k_m}})\}. \end{aligned}$$

Now (29) leads to

$$\lim_{m \rightarrow \infty} \frac{q'_{n_{k_m}}}{q_{n_{k_m}} + q'_{n_{k_m}}} = \alpha, \quad \lim_{m \rightarrow \infty} \frac{q_{n_{k_m}} + q'_{n_{k_m}}}{2q_{n_{k_m}} + q'_{n_{k_m}}} = \beta,$$

where by (25) and (26) one has $\beta = \frac{1}{\lambda}$, $\alpha = \frac{c}{\lambda}$ for $S(p_{n_{k_m}}/q_{n_{k_m}}) < S(p'_{n_{k_m}}/q'_{n_{k_m}})$, and $\alpha = \frac{c}{\lambda^2}$ for $S(p_{n_{k_m}}/q_{n_{k_m}}) > S(p'_{n_{k_m}}/q'_{n_{k_m}})$. Note that

$$\frac{2q_{n_{k_m}} + q'_{n_{k_m}}}{q_{n_{k_m}} + q'_{n_{k_m}}} = 2 - \frac{q'_{n_{k_m}}}{q_{n_{k_m}} + q'_{n_{k_m}}}.$$

So we have

$$\frac{1}{\beta} = 2 - \alpha.$$

For $\beta = \frac{1}{\lambda}$, $\alpha = \frac{c}{\lambda}$ we get

$$\lambda^2 - 2 = 0.$$

For $\beta = \frac{1}{\lambda}$, $\alpha = \frac{c}{\lambda^2}$ we get

$$\lambda^2 - \lambda - 1 = 0.$$

In both cases we have a contradiction with the fact that λ is a root of equation (13).

1,b) In this case

$$\begin{aligned} \{p_{n_{km}+1}/q_{n_{km}+1}, p'_{n_{km}+1}/q'_{n_{km}+1}\} &= \{p_{n_{km}}/q_{n_{km}}, (p_{n_{km}} + p'_{n_{km}})/(q_{n_{km}} + q'_{n_{km}})\}, \\ \{p_{n_{km}+2}/q_{n_{km}+2}, p'_{n_{km}+2}/q'_{n_{km}+2}\} &= \\ &= \{(2p_{n_{km}} + p'_{n_{km}})/(2q_{n_{km}} + q'_{n_{km}}), (p_{n_{km}} + p'_{n_{km}})/(q_{n_{km}} + q'_{n_{km}})\}, \end{aligned}$$

Now (29) leads to

$$\lim_{m \rightarrow \infty} \frac{q'_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}} = \alpha, \quad \lim_{m \rightarrow \infty} \frac{q_{n_{km}}}{2q_{n_{km}} + q'_{n_{km}}} = \beta,$$

where by (25) and (26) one has $\beta = \frac{1}{c\lambda}$, $\alpha = \frac{c}{\lambda}$ for $S(p_{n_{km}}/q_{n_{km}}) < S(p'_{n_{km}}/q'_{n_{km}})$, and $\alpha = \frac{c}{\lambda^2}$ for $S(p_{n_{km}}/q_{n_{km}}) > S(p'_{n_{km}}/q'_{n_{km}})$. Note that

$$\frac{2q_{n_{km}} + q'_{n_{km}}}{q_{n_{km}}} = 1 + \frac{q_{n_{km}} + q'_{n_{km}}}{q_{n_{km}}} = 1 + \frac{1}{1 - \frac{q'_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}}}.$$

So we have

$$\frac{1}{\beta} = 1 + \frac{1}{1 - \alpha}$$

For $\beta = \frac{1}{c\lambda}$, $\alpha = \frac{c}{\lambda}$ we get

$$\lambda^2 - 2 = 0.$$

For $\beta = \frac{1}{c\lambda}$, $\alpha = \frac{c}{\lambda^2}$

$$\lambda^2 - \lambda - 1 = 0.$$

Again in both cases we have the contradiction with the fact that λ is a root of equation (13).

2,a) In this case

$$\begin{aligned} \{p_{n_{km}+1}/q_{n_{km}+1}, p'_{n_{km}+1}/q'_{n_{km}+1}\} &= \{(p_{n_{km}} + p'_{n_{km}})/(q_{n_{km}} + q'_{n_{km}}), p'_{n_{km}}/q'_{n_{km}}\}, \\ \{p_{n_{km}+2}/q_{n_{km}+2}, p'_{n_{km}+2}/q'_{n_{km}+2}\} &= \\ &= \{(p_{n_{km}} + p'_{n_{km}})/(q_{n_{km}} + q'_{n_{km}}), (p_{n_{km}} + 2p'_{n_{km}})/(q_{n_{km}} + 2q'_{n_{km}})\}. \end{aligned}$$

So (29) leads to

$$\lim_{m \rightarrow \infty} \frac{q_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}} = \alpha, \quad \lim_{m \rightarrow \infty} \frac{q'_{n_{km}}}{q_{n_{km}} + 2q'_{n_{km}}} = \beta,$$

where by (27) and (28) one has $\beta = \frac{1}{c\lambda}$, $\alpha = \frac{c}{\lambda^2}$ for $S(p_{n_{km}}/q_{n_{km}}) < S(p'_{n_{km}}/q'_{n_{km}})$, and $\alpha = \frac{c}{\lambda}$ for $S(p_{n_{km}}/q_{n_{km}}) > S(p'_{n_{km}}/q'_{n_{km}})$. Note that

$$\frac{q_{n_{km}} + 2q'_{n_{km}}}{q'_{n_{km}}} = 1 + \frac{q_{n_{km}} + q'_{n_{km}}}{q'_{n_{km}}} = 1 + \frac{1}{1 - \frac{q_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}}}.$$

So we have

$$\frac{1}{\beta} = 1 + \frac{1}{1 - \alpha},$$

and tis case is reduced to the case 1,b).

2,b) In this case

$$\begin{aligned} \{p_{n_{km}+1}/q_{n_{km}+1}, p'_{n_{km}+1}/q'_{n_{km}+1}\} &= \{(p_{n_{km}} + p'_{n_{km}})/(q_{n_{km}} + q'_{n_{km}}), p'_{n_{km}}/q'_{n_{km}}\}, \\ \{p_{n_{km}+2}/q_{n_{km}+2}, p'_{n_{km}+2}/q'_{n_{km}+2}\} &= \{(p_{n_{km}} + 2p'_{n_{km}})/(q_{n_{km}} + 2q'_{n_{km}}), p'_{n_{km}}/q'_{n_{km}}\}. \end{aligned}$$

Now (29) leads to

$$\lim_{m \rightarrow \infty} \frac{q_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}} = \alpha, \quad \lim_{m \rightarrow \infty} \frac{q_{n_{km}} + q'_{n_{km}}}{q_{n_{km}} + 2q'_{n_{km}}} = \beta,$$

where by (27) and (28) one has $\beta = \frac{1}{\lambda}$, $\alpha = \frac{c}{\lambda^2}$ for $S(p_{n_{km}}/q_{n_{km}}) < S(p'_{n_{km}}/q'_{n_{km}})$, and $\alpha = \frac{c}{\lambda}$ for $S(p_{n_{km}}/q_{n_{km}}) > S(p'_{n_{km}}/q'_{n_{km}})$. Note that

$$\frac{q_{n_{km}} + 2q'_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}} = 2 - \frac{q_{n_{km}}}{q_{n_{km}} + q'_{n_{km}}}.$$

So we have

$$\frac{1}{\beta} = 2 - \alpha,$$

and we have the same situation as in the case 1,a).

Theorem 2 is proved. ■

References

- [1] A. Denjoy. Sur une fonction reele de Minkowski. — J. Math. Pures Appl. 17(1938), pp. 105-151.

- [2] A. A. Dushistova, I. D. Kan and N. G. Moshchevitin. — Differentiability of the Minkowski question mark function. Preprint available at arXiv:0903.5537v1 (2009)
- [3] Yu. Yu Finkel'shtein. Klein polygons and reduced regular continued fractions. — Russian Mathematical Surveys (1993), 48(3):198.
- [4] E. N. Jabitskaya. On arithmetical nature of Tichy-Uitz's function. — To appear in *Functiones et Approximatio*. Preprint available at arXiv:0909.1273v2 (2009)
- [5] E. N. Zhabitskaya. The average length of reduced regular continued fractions. — *Sbornik: Mathematics*, 2009, 200:8, 1181-1214.
- [6] O. Perron. *Die Lehre von den Kettenbrüchen*. — Bd. I. Teuber, 1954.
- [7] R. Salem. On some singular monotonic functions which are strictly increasing. — *Trans. Amer. Math. Soc.* 53(1943), pp. 427-439.
- [8] R. F. Tichy, J. Uitz. An extension of Minkowski's singular function. — *Appl. Math. Lett.*, 8 (1995), 39-46.
- [9] A. V. Ustinov. Asymptotic behaviour of the first and second moments for the number of steps in the Euclidean algorithm. — *Izvestiya: Mathematics*, 2008, 72:5, 1023-1059.
- [10] A. V. Ustinov. The mean number of steps in the Euclidean algorithm with least absolute value remainders. — *Mathematical Notes*, 2009, 85:1, 142-145.
- [11] B. Vallee. A unifying framework for the analysis of class of Euclidean algorithms. — *Proceedings of LATIN'2000, Lecture Notes in Computer Science 1776*, Springer, 343-354.
- [12] B. Vallee. Dynamical analysis of a class of Euclidean algorithms. — *Theoretical computer science*, v. 297/1-3, 2003, 447-486.
- [13] B. Vallee, V. Baladi. Euclidean algorithms are Gaussian. — *J. Number Theory*, v. 110, 2005, 331-386.
- [14] P. Viader, J. Paradis, L. Bibiloni. A new light of Minkowski's $?(x)$ function. — *J. Number Theory.*, 73 (1998), 212 -227.